

Stereo sensitivity of exchange interactions in Ni^{II} and Cu^{II} heterospin complexes with 5-formylpyrrolyl-substituted nitroxides

Zueva E., Tretyakov E., Fokin S., Romanenko G., Tkacheva A., Bogomyakov A., Petrova O., Trofimov B., Sagdeev R., Ovcharenko V.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Science+Business Media New York. 5-Formylpyrrolyl-substituted nitronyl and imino nitroxide radicals HL1 and HL2 were synthesized. Their solid phases are formed by packing pairs of the molecules. In the {HL1..HL1} pairs, the dominant interaction is the ferromagnetic exchange with $J/kB = 8.8$ K (Hamiltonian $H = 2 J(s_1 \rightarrow \cdot s_2 \rightarrow)$). The ferromagnetic exchange occurs also in the heterospin molecules [Ni(L1)2], [Cu(L1)2], and [Ni(L2)2(MeOH)2]. In the complexes [Ni(L1)2] and [Cu(L1)2], a small change in the mutual orientation of the coordinated ligands has a considerable effect on the value and the sign of the energy of exchange interactions between the unpaired electrons of the metal ion and paramagnetic ligands.

<http://dx.doi.org/10.1007/s11172-016-1353-8>

Keywords

copper(II) complexes, nickel(II) complexes, nitroxide radicals, pyrrole, quantum chemical calculations, X-ray diffraction